

SMILE



GIT HISTOLOGY IN PICTURES

BY: **AM**

Accessory digestive organs

Parotid salivary gland
Teeth
Tongue
Sublingual salivary gland
Submandibular salivary gland

**Gastrointestinal tract
(digestive organs)**

Oral cavity
Pharynx

Esophagus

Liver


Stomach

Gallbladder
Pancreas

Large intestine


Small intestine

Anus



THE LIP





OM : oral mucosa


G : minor salivary glands

M : striated muscle

F : hair follicles


S : skin


THE LIP




The **lips** are the entry point of the alimentary canal. Here, the thin **keratinized epithelium** of face skin changes to the thick **parakeratinized epithelium** of the oral mucosa. At the mucocutaneous junction, the red portion of the lips, is characterized by deep penetration of connective tissue papillae into the base of the **stratified squamous keratinized epithelium**. The blood vessels and nerve endings in these papillae are responsible for both the color and the exquisite touch sensitivity of the lips.

ORIENTATION MICROGRAPH: An H&E-stained sagittal section through the upper lip in this low-power orientation photomicrograph to the right ($\times 8$) reveals the skin of the face, the red margin of the lip, and the transition to the oral mucosa (OM). The *marked rectangles* indicate representative areas of each of these sites, shown at higher magnifications in upper, middle and lower rows of figures, on the adjacent plate. Note the change in thickness of the epithelium from the exterior or facial portion of the lip (the vertical surface on the right) to the interior surface of the oral cavity (the surface beginning with *rectangle marked lower* and continuing down the left surface of the lip) in this micrograph.






△ Early carcinoma of the lip.



▲ LMs of parts of the lip. **Left**, The vermillion border is stratified squamous epithelium (SSE) with a thin layer of surface keratin, below. Underlying connective tissue—lamina propria (LP)—contains many blood vessels (BV). The highly corrugated interface between epithelium and connective tissue shows tall papillae (*) penetrating the epithelium to take capillaries close to the surface. **Right**, The external cutaneous surface, of typical thin skin, consists of epidermis (Ep) and underlying dermis (De). A hair follicle (HF) and associated hair shaft (HS) are seen. **Left**: 130×; **Right**: 85×. H&E.


THE TONGUE

Three parallel white lines of varying lengths and slopes are positioned in the bottom right corner of the image, extending from the right edge towards the center.



Epithelium


Filiform papilla



Epithelium

Taste bud

Fungiform papilla




Root of tongue

Body of tongue

Apex of tongue


Dorsal surface of tongue



Epithelium

Taste bud


Vallate papilla





Epithelium




Taste bud

Foliate papilla





TASTE BUDS






FIGURE 13.2 ■ Anterior region of the tongue: apex (longitudinal section). Stain: hematoxylin and eosin. Low magnification.



FIGURE 13.3 ■ Tongue: circumvallate papilla (cross section). Stain: hematoxylin and eosin. Medium magnification.




FIGURE 13.4 ■ Tongue: filiform and fungiform papillae. Stain: hematoxylin and eosin. $\times 25$.






FIGURE 13.5 ■ Tongue: taste buds. Stain: hematoxylin and eosin. High magnification.

THE ALIMENTARY CANAL


Three parallel white lines of varying lengths are positioned diagonally on the right side of the image, extending from the middle of the word 'ALIMENTARY' down towards the bottom right corner.

Major layers and organization of the digestive tract





THE ESOPHAGUS




a



(a) In cross section the four major layers of the GI tract are clearly seen. The esophageal mucosa is folded longitudinally, with the lumen largely closed. X10. H&E. **(b)** Higher magnification of

the mucosa shows the stratified squamous epithelium (**E**), the lamina propria (**LP**) with scattered lymphocytes, and strands of smooth muscle in the muscularis mucosae (**MM**). X65. H&E.




Stratified squamous
epithelium

Lamina
propria


Smooth
muscle

Esophageal
glands


Skeletal
muscle



Photomicrograph
of a section of the
upper region of the
esophagus ...



Photomicrograph of the esophageal mucosa



Photomicrograph of an esophageal submucosal gland




FIGURE 14.1 ■ Wall of the upper esophagus (transverse section). Stain: hematoxylin and eosin. Low magnification.



FIGURE 14.2 ■ Upper esophagus (transverse section). Stain: hematoxylin and eosin. Low magnification.




FIGURE 14.3 ■ Lower esophagus (transverse section). Stain: hematoxylin and eosin. Low magnification.




FIGURE 14.4 ■ Upper esophagus: mucosa and submucosa (longitudinal view). Stain: hematoxylin and eosin. Medium magnification.









FIGURE 14.5 ■ Lower esophageal wall (transverse section). Stain: Mallory-Azan. ×30.

THE STOMACH








© Elsevier. Gartner & Hiatt: Color Textbook of Histology 3E - www.studentconsult.com

[Add to My Slides](#) [Go to My Slides](#)

Figure 17-3 Cellular composition of the fundic stomach and fundic gland. The fundic glands open into the bottom of the gastric pits, and each gland is subdivided into an isthmus, a neck, and a base.



Stomach and cells







FIGURE 17.14 • Photomicrograph of cardiac glands.



FIGURE 17.15 • Photomicrograph of pyloric glands.





Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Composite diagram of a parietal cell, showing the ultrastructural differences between a resting cell (**left**) and an active cell (**right**). Note that the tubulovesicles (TV) in the cytoplasm of the resting cell fuse to form microvilli (MV) that fill up the intracellular canaliculi (IC). G, Golgi complex; M, mitochondria. (Based on the work of Ito S, Schofield GC: Studies on the depletion and accumulation of microvilli and changes in the tubulovesicular compartment of mouse parietal cells in relation to gastric acid secretion. *J Cell Biol* 1974;63:364.)






FIGURE 17.12 • Electron micrograph and diagrams of enteroendocrine cells. **a.** This electron micrograph shows an example of the "closed" enteroendocrine cell. *Arrowheads* mark the boundary between the enteroendocrine cell and the adjacent epithelial cells. At its base, the enteroendocrine cell rests on the basal lamina (BL). This cell does not extend to the epithelial or luminal surface. Numerous secretory vesicle (G) in the base of the cell are secreted in the direction of the *arrows* across the basal lamina and into the connective tissue (CT). *En*, endothelium of capillary; *M*, mitochondria; *rER*, rough endoplasmic reticulum; *sER*, smooth endoplasmic reticulum. **b.** This diagram of an enteroendocrine "closed" cell is drawn to show that it does not reach the epithelial surface. The secretory vesicles are regularly lost during routine preparation. Because of the absence of other distinctive organelles, the nucleus appears to be surrounded by a small amount of clear cytoplasm in H&E-stained sections. **c.** The enteroendocrine "open" extend to the epithelial surface. Microvilli on the apical surface of these cells possess taste receptors and are able to detect sweet, bitter, and umami sensations. These cells serve as chemoreceptor cells, which monitor an environment on the surface of the epithelium. They are involved in a regulation of gastrointestinal hormones secretion.

THE SMALL INTESTINE

Three parallel white lines of varying lengths are positioned diagonally in the bottom right corner of the image.










FIG. 14.18 Small intestine



Source: Mescher AL: *Junqueira's Basic Histology: Text and Atlas*, 12th Edition: <http://www.accessmedicine.com>
Copyright © The McGraw-Hill Companies, Inc. All rights reserved.

Plicae circulares (folds) of the jejunum.

The mucosa and submucosa (SM) of the small intestine form distinct projecting folds called plicae (P), which encircle or spiral around the inner circumference and are best developed in the jejunum. On each fold the mucosa forms a dense covering of projecting structures called villi (V). In this longitudinal section the two layers of the muscularis (M) are clearly distinguished. The inner layer has smooth muscle encircling the submucosa; the outer layer runs lengthwise just inside the serosa (S), the gut's outer layer. This arrangement of smooth muscle provides for strong peristaltic movement of the gut's contents. X12. H&E.



© Elsevier. Gartner & Hiatt: Color Textbook of Histology 3E - www.studentconsult.com

[Add to MySlides](#) [Go to MySlides](#)

Figure 17-13 Mucosa, villi, crypts of Lieberkühn, and component cells of the small intestine. Note that the crypts of Lieberkühn open into the intervillar spaces. There is a solitary lymphoid nodule in the lamina propria.





FIGURE 15.4 ■ Intestinal glands with Paneth cells and enteroendocrine cells. Stain: hematoxylin and eosin. High magnification.



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.


Photomicrograph of the small intestine. Note the villi, intestinal glands, submucosa, muscle layers, and serosa. PT stain. Low magnification.


C



intestinal villus

A





Source: Mescher AL: *Junqueira's Basic Histology: Text and Atlas, 12th Edition*: <http://www.accessmedicine.com>

Copyright © The McGraw-Hill Companies, Inc. All rights reserved.

Absorptive surface of the small intestine.




FIGURE 15.7 ■ Small intestine: villi (longitudinal and transverse sections). Stain: periodic acid-Schiff. Medium magnification.





FIGURE 15.8 ■ Ultrastructure of microvilli in an absorptive cell in the small intestine. Courtesy of Dr. Rex A. Hess, Professor Emeritus, Comparative Biosciences, College of Veterinary Medicine, University of Illinois, Urbana, Illinois. $\times 6,150$.




Source: Mescher AL: *Junqueira's Basic Histology: Text and Atlas*, 12th Edition: <http://www.accessmedicine.com>


Copyright © The McGraw-Hill Companies, Inc. All rights reserved.

Microvasculature, lymphatics, and muscle in villi.

Stomach




Small intestine



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Regeneration of the epithelial lining of the stomach and small intestine. Note differences in the location of stem cells.



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Photomicrograph of the duodenum, showing villi and duodenal glands in the submucosa. H&E stain. Low magnification. (Courtesy of MF Santos.)




FIGURE 15.2 ■ Small intestine: duodenum (transverse section). Stain: hematoxylin and eosin. x25.




FIGURE 15.1 ■ Small intestine: duodenum (longitudinal section). Stain: hematoxylin and eosin. Low magnification.




FIGURE 15.3 ■ Small intestine: jejunum (transverse section). Stain: hematoxylin and eosin. Low magnification.




FIGURE 15.5 ■ Small intestine: jejunum with Paneth cells. Stain: Mallory-Azan. $\times 40$.




FIGURE 15.6 ■ Small intestine: ileum with lymphatic nodules (Peyer patches) (transverse section). Stain: hematoxylin and eosin. Low magnification.

DON'T GIVE UP



THE LARGE INTESTINE

Three parallel white diagonal lines are located on the right side of the image, extending from the middle towards the bottom right corner.






FIGURE 15.9 ■ Large intestine: colon and mesentery (panoramic view, transverse section).
Stain: hematoxylin and eosin. Low magnification.



LARGE INTESTINE



G = Goblet cells



CL = Crypts of Lieberkuhn



LP = Lamina propria

SM = Submucosa


MM = Muscularis mucosa

O = Open lumen of crypts of Lieberkuhn






Transverse section of the colon shows the muscularis externa (ME), including a **tenia coli** cut transversely in the lower part of the figure, the submucosa (S), the mucosa (M) filled with **tubular intestinal glands**. Some of these glands are cut longitudinally, but most seen here are cut transversely.



Transversely cut glands are seen to consist of simple columnar epithelium surrounded by a tubular lumen (L) and embedded in lamina propria (LP) with many free lymphocytes. Lymphocytes can also be seen penetrating the epithelium (arrow)



Longitudinal section of one intestinal gland stained for glycoproteins shows mucus in the lumen and two major cell types in the epithelium: goblet cells (G) and the neighboring columnar cells specialized for water absorption.






FIGURE 15.10 ■ Large intestine: colon wall (transverse section). Stain: hematoxylin and eosin. x30.



▲ **LM of the colonic mucosa.** Surface epithelium containing goblet cells (G) and enterocytes (En) invaginates to form an intestinal crypt. The lamina propria (LP), with capillaries (arrows) and larger blood vessels (BV), is richly cellular. 600 \times . Toluidine blue.



EM of the colonic mucosa. Enterocytes have an apical striated border of microvilli that project into the lumen.

These cells occur with goblet cells, which have apical mucin granules. The lamina propria contains a capillary (Cap).







FIGURE 15.11 ■ Large intestine: colon wall (transverse section). Stain: hematoxylin and eosin. Medium magnification.




▼ Ileocecal region.



▼ Cross section of appendix.





THE VERMIFORM APPENDIX






FIGURE 15.12 ■ Appendix (panoramic view, transverse section). Stain: hematoxylin and eosin. Low magnification.





▲ **LM of the appendix in transverse section.**

Prominent mucosa (**Mu**) shows multiple lymphoid nodules (**LN**) bulging into a narrow stellate lumen (*). Connective tissue makes up a submucosa (**SM**); the muscularis externa (**ME**) contains two layers of smooth muscle. 8×. H&E.




▲ **LM of part of the appendix.** Mucosa contains richly cellular lamina propria (**LP**) infiltrated with lymphoid nodules (**LN**) that extend into submucosa (**SM**). Shallow invaginations of the surface—intestinal crypts (**arrows**)—vary in length. The submucosa has some adipose tissue. 85×. H&E.




DRAWING OF THE RECTUM AND ANAL CANAL


▼ Rectum and anal canal.




F. Netter M.D.



◀ **Endoscopic view of the anorectal junction.** The transition from rectum to anal canal is near the serrated pectinate line (arrows), which shows a color change on the surface. It marks the squamocolumnar junction.



▲ **LM of the anorectal junction in longitudinal section.** The rectum (at top) leads into the anal canal (at bottom). The squamocolumnar junction occurs well above the pectinate line. Other features are indicated. Superior (SH) and inferior (IH) hemorrhoidal venous plexuses—thin-walled veins filled with blood—lie on both sides of the pectinate line. 5×. H&E.



LM OF THE MUCOSA OF THE ANAL CANAL

Further along the canal, epithelium becomes **stratified squamous**. Just below the pectinate line, it is **nonkeratinized**, but it becomes keratinized. In connective tissue under the epithelium is a plexus of **thin-walled hemorrhoidal veins**.

THE SITE OF
JUNCTION BETWEEN
THE DIFFERENT
PARTS OF THE
DIGESTIVE TUBE




FIGURE 14.6 ■ Esophageal–stomach junction. Stain: hematoxylin and eosin. Low magnification.




FIGURE 14.14 ■ Pyloric-duodenal junction (longitudinal section). Stain: hematoxylin and eosin. Low magnification.






FIGURE 15.14 ■ Anorectal junction (longitudinal section). Stain: hematoxylin and eosin. Low magnification.




LMs of the anorectal junction at low (Left) and high (Right) magnifications


An abrupt transition (**double arrows**) from simple columnar (left) to stratified columnar (right) epithelium occurs in this part of the pectinate line. Mucus-secreting goblet cells predominate on the mucosal surface and in rectal crypts. Highly cellular lamina propria under the epithelium contains lymphoid nodules.

THE SALIVARY GLANDS






MAJOR SALIVARY GLANDS



SALIVARY SECRETORY UNIT



The secretory portions are composed of pyramidal serous (violet) and mucous (tan) cells. **Serous cells** are typical protein-secreting cells, with rounded nuclei, accumulation of rough ER in the basal third, and an apex filled with protein-rich secretory granules. The nuclei of **mucous cells**, flattened with condensed chromatin, are located near the bases of the cells.

The short intercalated ducts are lined with cuboidal epithelium. The **striated ducts** are composed of columnar cells with characteristics of ion-transporting cells: basal membrane invaginations with mitochondrial accumulations. **Myoepithelial cells** are shown around the serous acini.




FIGURE 13.16 ■ Serous salivary gland: parotid gland. Stain: hematoxylin and eosin. $\times 165$.




FIGURE 13.17 ■ Mixed salivary gland: sublingual gland. Stain: hematoxylin and eosin. $\times 165$.




FIG. 13.17 Striated ducts
H&E (HP)

The striated ducts **D** are lined by tall columnar cells with large nuclei located towards the apex of the cell. The basal cytoplasm appears striated, reflecting the presence of basal interdigitations of cytoplasmic processes of adjacent cells and associated columns of mitochondria. This feature greatly extends the area of membrane available for exchange of water and ions, in a similar fashion to the proximal convoluted tubule of the kidney (see [Fig. 16.17](#)). The duct epithelium also secretes lysozyme and immunoglobulin (Ig)A. In predominantly serous salivary glands, the striated ducts are larger than in predominantly mucous glands, a feature associated with the role of the striated duct in modifying isotonic basic saliva to produce hypotonic saliva. The sparse supporting tissue between the secretory acini contains a rich network of capillaries **C**.




FIGURE 13.13 ■ Parotid salivary gland. Stain: hematoxylin and eosin. Upper: medium magnification. Lower: high magnification.




FIGURE 13.14 ■ Submandibular salivary gland. Stain: hematoxylin and eosin. Upper: medium magnification. Lower: high magnification.






FIGURE 13.15 ■ Sublingual salivary gland. Stain: hematoxylin and eosin. Upper: medium magnification. Lower: high magnification.




The large parotid gland consists entirely of serous acini with cells producing amylase and other proteins for storage in secretory granules. **(a)** Micrograph of a parotid gland shows densely packed serous acini (A) with ducts. Secretory granules of serous cells are clearly shown in this plastic section,



as well as an intercalated duct (ID) and striated duct (SD), both cut transversely. X400. PT.

(b) Striations of a duct (SD) are better seen here, along with a septum (CT) and numerous serous acini (A). The connective tissue often includes adipocytes. X200. H&E.


FIGURE 16–4 Ultrastructure of serous and mucous cells.




A micrograph of a mixed acinus from a submandibular gland shows both serous and mucous cells surrounding the small lumen (L). Mucous cells (M) have large, hydrophilic granules like those of goblet cells, while serous cells (S) have small, dense granules. Small myoepithelial cells (My) extend contractile processes around each acinus. X2500.

(With permission, from Dr John D. Harrison, King's College London Dental Institute, London, UK.)

FIGURE 16-5 Submandibular gland and sublingual gland.




(a) The submandibular gland is a mixed serous and mucous gland (serous cells predominate), and shows well-stained serous acini (**A**) and serous demilunes (**S**) and pale-staining mucous cells (**M**) grouped as tubules in this tubuloacinar gland. Small intralobular ducts (**ID**) drain each lobule. X340. H&E.



(b) The sublingual gland is a mixed but largely mucous gland with a tubuloacinar arrangement of poorly stained mucous cells (**M**). Small intralobular ducts (**ID**) are seen in connective tissue, as well as small fascicles of lingual striated muscle (**SM**). X140. H&E.

THE PANCREAS

Three parallel white lines of increasing length, slanted upwards from left to right, located in the bottom right corner of the image.










FIGURE 16-9 Pancreatic acini.




(a) Micrograph of exocrine pancreas shows the serous, enzyme-producing cells arranged in small acini (**A**) with very small lumens. Acini are surrounded by only small amounts of connective tissue with fibroblasts (**F**). Each acinus is drained by an intercalated duct with its initial cells, the centroacinar cells (**arrow**), inserted into the acinar lumen. X200. H&E.



(b) The diagram shows the arrangement of cells more clearly. Under the influence of secretin, the **centroacinar** and **intercalated duct cells** secrete a copious HCO_3^- -rich fluid that hydrates, flushes, and alkalinizes the enzymatic secretion of the acini.




PANCREATIC ACINUS AND ITS DUCT SYSTEM




Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.



Section of the exocrine pancreas showing its main components. PT stain. Medium magnification.




Electron micrograph of the pancreatic acinus and intercalated duct



◀ **LM of the pancreas at low magnification.** Intensely stained parenchyma of the gland is organized into lobules. More lightly stained stroma consists mainly of connective tissue. Thumbnail areas in the three boxes are seen at higher magnification below, and show different parts of the excretory duct system traversing the gland. 18×. H&E.



▲ ► **LMs of the pancreas showing parts of the exocrine duct system.** Short, branching intercalated ducts—the initial part of the excretory duct system—drain acini and are lined by a low simple cuboidal epithelium (A). Longer intralobular ducts are lined by simple cuboidal epithelium consisting of more plump cells (B). Larger interlobular ducts are lined by simple cuboidal epithelium invested by dense fibrous connective tissue (C). Interlobular ducts, in turn, drain directly into the main pancreatic ducts, which deliver secretions to the duodenum. Secretory cells of acini are intensely eosinophilic, have basal nuclei, and produce digestive enzymes. Centroacinar cells protrude into the lumen of each acinus. They have clear cytoplasm, and secrete fluid rich in HCO_3^- and Na^+ . Pale cells in the richly vascularized islets constitute the endocrine part of the gland. **A:** 290×; **B:** 215×; **C:** 250×. H&E.




DON'T GIVE
UP


YOU ARE CLOSER THAN YOU THINK

#FUCOS_ON_YOUR_FUTURE ☺

THE

LIVER






© Elsevier. Gartner & Hiatt: Color Textbook of Histology 3E - www.studentconsult.com

[Add to My Slides](#) [Go to My Slides](#)

Figure 18-9 Liver. A, Gross anatomy of the liver. B, Liver lobules displaying the portal areas and the central vein. C, Portion of the liver lobule displaying the portal area, liver plates, sinusoids, and bile canaliculi.



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Three-dimensional aspect of the normal liver. In the upper center is the central vein; in the lower center is the portal vein. Note the bile canaliculus, liver plates, Hering's canal, Kupffer cells, sinusoid, fat-storing cell, and sinusoid endothelial cells. (Courtesy of M Muto.)




FIGURE 16.8 ■ Reticular fibers in liver lobule. Stain: reticulin method. Medium magnification.






FIGURE 16.1 ■ Pig liver (panoramic view, transverse section). Stain: Mallory-Azan. Low magnification.



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Ultrastructure of a hepatocyte. RER, rough endoplasmic reticulum; SER, smooth endoplasmic reticulum. $\times 10,000$.




© Elsevier. Gartner & Hiatt: Color Textbook of Histology 3E - www.studentconsult.com



[Add to My Slides](#) [Go to My Slides](#)



Figure 18-14 A hepatocyte and its sinusoidal and lateral domains. ER, endoplasmic reticulum. (From Lentz TL: Cell Fine Structure: An Atlas of Drawings of Whole-Cell Structure. Philadelphia, WB Saunders, 1971.)

Relationship Between Hepatocytes and Sinusoids

© S.C. Kempf 2/2000 Histology 0509













Diagram of a classic liver lobule.



CLASSIC LOBULE



PORTAL LOBULE



LIVER ACINUS






FIGURE 18.6 • The liver acinus. The liver acinus is a functional interpretation of liver organization. It consists of adjacent sectors of neighboring hexagonal fields of classic lobules partially separated by distributing blood vessels. The zones, marked 1, 2, and 3, are supplied with blood that is most oxygenated and richest in nutrients in zone 1 and least so in zone 3. The terminal hepatic venules (central veins) in this interpretation are at the edges of the acinus instead of in the center, as in the classic lobule. The vessels of the portal canals, namely, terminal branches of the portal vein and hepatic artery that, along with the smallest bile ducts, make up the portal triad, are shown at the corners of the hexagon that outlines the cross-sectioned profile of the classic lobule.




NB;


- the bile flows **away from** the central vein



NB;

- the blood flows **towards** the central vein











Diagram of the flow of blood and bile in the liver



BLOOD SUPPLY TO THE LIVER: THE PORTAL TRIAD

GALL BLADDER










FIGURE 16.15 ■ Wall of the gallbladder. Stain: hematoxylin and eosin. Low magnification.



The gallbladder is a saclike structure that stores and concentrates bile, and releases it into the duodenum after a meal. **(a)** Its wall consists largely of a highly folded mucosa, with a simple columnar epithelium (arrows) overlying a typical lamina propria (**LP**); a muscularis (**M**) with bundles of muscle fibers oriented in all directions to facilitate emptying of the organ; and an external adventitia (**A**) where it is against the liver and a serosa where it is exposed. X60. H&E.















(b) TEM of the epithelium shows cells specialized for water uptake across apical microvilli (**MV**) and release into the intercellular spaces (**arrows**) along the folded basolateral cell membranes. From these spaces water is quickly removed by capillaries in the lamina propria. Abundant mitochondria provide the energy for this pumping process. Scattered apical secretory granules (**G**) contain mucus. X5600.



FIGURE 18.16 • Photomicrograph of the wall of the gallbladder. The mucosa of the gallbladder consists of a lining of simple columnar epithelial cells and a lamina propria of loose connective tissue, which typically exhibits numerous deep folds in the mucosa. Beneath this layer is a relatively thick layer, the muscularis externa. There is no muscularis mucosae or submucosa. The smooth muscle bundles of the muscularis externa are randomly oriented. External to the muscle is an adventitia containing adipose tissue and blood vessels. The portion of the gallbladder not attached to the liver displays a typical serosa instead of an adventitia. $\times 175$.

REFERENCES

 Color Atlas and Text of Histology 2013 6th	12/26/2014 3:39 PM	Adobe Acrobat D...	97,158 KB
 diFiore's Atlas of Histology with Function...	1/13/2015 12:39 PM	Adobe Acrobat D...	427,267 KB
 Histology A Text and Atlas, with Correlat...	3/18/2012 2:43 PM	Adobe Acrobat D...	146,479 KB
 Junqueira's histology 2013 13th	12/26/2014 3:01 PM	Adobe Acrobat D...	115,372 KB
 Netter's Essential Histology 2nd 2013	1/16/2015 6:44 AM	Adobe Acrobat D...	452,517 KB
 Wheater's Functional Histology, A Text a...	12/25/2014 11:14 ...	Adobe Acrobat D...	245,057 KB
 Basic Histology ASM2016 Virtual Library	3/18/2007 1:59 AM	Compiled HTML ...	78,581 KB
 Color Atlas of Cytology, Histology, and ...	2/9/2007 9:33 PM	Adobe Acrobat D...	25,189 KB
 Color Textbook of Histology, Hiatt ASM2...	4/4/2011 10:55 PM	Compiled HTML ...	165,984 KB
 Essential Histology, 2nd ed. ASM2016 Virt...	7/14/2007 8:16 PM	Adobe Acrobat D...	110,262 KB
 Junqueira's Basic Histology ASM2016 Virt...	2/23/2010 8:10 AM	Compiled HTML ...	130,619 KB
 Kuehnel, Color Atlas of Cytology, Histolo...	2/10/2007 4:16 AM	Adobe Acrobat D...	112,755 KB

“REMEMBER”

YOU CAN MAKE
A DIFFERENCE

BY: ALI

MAHMOUD